

## Antimikrobielle Wirkung von ätherischen Ölen

### Literatur

- Aruoma OI (1998). Free radicals, oxidative stress, and antioxidants in human health and disease. *Journal of the American Oil Chemist's Society* 75: 199–212.
- Bakkeren E et al. (2020). Evolutionary causes and consequences of bacterial antibiotic persistences. *Nature reviews Microbiology* 18: 479–490.
- Benali T et al. (2021). Phytochemical analysis and study of antioxidant, anticandidal, and antibacterial activities of *Teucrium polium* subsp. *polium* and *Micromeria graeca* (Lamiaceae) essential oils from Northern Morocco. *Evidence-Based Complementary and Alternative Medicine*: 6641720.
- Bezerra Filho CM et al. (2020). Antimicrobial and antivirulence action of *Eugenia brejoensis* essential oil in vitro and in vivo invertebrate models. *Frontiers in Microbiology* 11: 424.
- Bogavac M et al. (2015). Alternative treatment of vaginal infections – in vitro antimicrobial and toxic effects of *Coriandrum sativum* L. and *Thymus vulgaris* L. essential oils. *Journal of Applied Microbiology* 119: 697–710.
- Böhm M (2000). Randomisierte, klinische Vergleichsstudie zur Behandlung eines Destillates ätherischer Öle in der chirurgischen Wundbehandlung. Dissertation an der Medizinischen Fakultät der Ludwig-Maximilian-Universität München.
- Boskovic M et al. (2015). Antimicrobial activity of thyme (*Thymus vulgaris*) and oregano (*Origanum vulgare*) essential oils against some food-borne microorganisms. *Procedia Food Science* 5: 18–21.
- Brun P et al. (2019). In vitro antimicrobial activities of commercially available tea tree (*Melaleuca alternifolia*) essential oils. *Current Microbiology* 76: 108–116.
- Burt S (2004). Essential oils: Their antibacterial properties and potential applications in foods. *International Journal of Food Microbiology* 94: 223–253.
- Caelli M et al. (2000). Tea tree oil as an alternative topical decolonization agent for methicillin-resistant *Staphylococcus aureus*. *Journal of Hospital Infection* 46: 236–237.
- Carmo ES et al. (2013). Treatment of pityriasis versicolor with topical application of essential oil of *Cymbopogon citratus* (DC) Stapf – therapeutic pilot study. *Anais Brasileiros de Dermatologia* 88: 381–385.
- Chaisripipat W et al. (2015). Anti-dandruff hair tonic containing lemongrass (*Cymbopogon flexuosus*) oil. *Forschende Komplementärmedizin* 22: 226–229.
- Chao S et al. (2008). Inhibition of methicillin-resistant *Staphylococcus aureus* (MRSA) by essential oils. *Flavour and Fragrance Journal* 23: 444–449.
- Cimanga K et al. (2002). Correlation between chemical composition and antibacterial activity of essential oils of some aromatic medicinal plants growing in the Democratic Republic of Congo. *Journal of Ethnopharmacology* 79: 213–220.
- Cowan MM (1999). Plant products as antimicrobial agents. *Clinical Microbiology Reviews* 12: 564–582.
- Cutillas AB et al. (2017). *Salvia officinalis* L. essential oils from Spain: determination of composition, antioxidant capacity, antienzymatic, and antimicrobial bioactivities. *Chemistry & Biodiversity*: 14.
- Davidson PM et al. (1997). Chemical preservations and natural antimicrobial compounds. In: Doyle et al. *Food Microbiology: Fundamentals and Frontiers*. Washington DC: AMS Press: 520–556.
- Diard M, Hardt WD (2017). Evolution of bacterial virulence. *FEMS Microbiology Reviews* 41: 679–697.
- Dorman HJD, Deans SG (2000). Antimicrobial agents from plants: Antibacterial activity of plant volatile oils. *Journal of Applied Microbiology* 88: 308–316.
- Dryden MS et al. (2004). A randomized, controlled trial of tea tree topical preparations versus a standard topical regimen for the clearance of MRSA colonization. *Journal of Hospital Infection* 56: 283–286.
- Feyaerts AF et al. (2018). Essential oils and their components are a class of antifungals with potent vapour-phase-mediated anti-*Candida* activity. *Scientific Report* 8: 3958.
- Fikry S et al. (2019). Chemical profiling, biostatic and biocidal dynamics of *Origanum vulgare* L. essential oil. *AMB Express* 9: 41.
- Fu Y et al. (2007). Antimicrobial activity of clove and rosemary essential oils alone and in combination. *Phytotherapy Research* 21: 989–994.
- Gaspar de Toledo L et al. (2020). Improved in vitro and in vivo anti-*Candida albicans* activity of *Cymbopogon nardus* essential oil by its incorporation into a microemulsion system. *International Journal of Nanomedicine* 15: 10481–10497.
- Gutierrez J et al. (2008). The antimicrobial efficacy of plant essential oils combinations and interactions with food ingredients. *International Journal of Food Microbiology* 124: 91–97.
- Hammer KA, Carson CF (2011). Chapter 11 – Antibacterial and antifungal activities of essential oils. In: Thormar H. *Lipids and essential oils as antimicrobial agents*. Chichester: John Wiley & Sons; 255–306.
- Harmati M et al. (2017). Binary mixture of *Satureja hortensis* and *Origanum vulgare* subsp. *hirtum* essential oils: in vivo therapeutic efficiency against *Helicobacter pylori* infection. *Helicobacter* 22: e12350.
- Hemaiswarya S et al. (2008). Synergism between natural products and antibiotics against infectious diseases. *Phytomedicine* 15: 639–652.
- Hendry ER et al. (2009). Antimicrobial efficacy of eucalyptus oil and 1,8-cineole alone and in combination with chlorhexidine digluconate against microorganisms grown in planktonic and biofilm cultures. *Journal of Antimicrobial Chemotherapy* 64: 1219–1225.
- Hu W et al. (2019). Antibacterial activity and mechanism of *Litsea cubeba* essential oil against methicillin-resistant *Staphylococcus aureus* (MRSA). *Industrial Crops and Products* 130: 34–41.
- Hulin V et al. (1998). Les propriétés antimicrobiennes des huiles essentielles et composés d'arômes. *Science des aliments* 18: 563–582.
- Karpanen TJ et al. (2008). Antimicrobial efficacy of chlorhexidine digluconate alone and in combination with eucalyptus oil, tea tree oil and thymol against planktonic and biofilm cultures of *Staphylococcus epidermidis*. *Journal of Antimicrobial Chemotherapy* 62: 1031–1036.
- Kelly D (1998). The physiology and metabolism of the human gastric pathogen (*Helicobacter pylori*). *Advances in Microbial Physiology* 40: 137–189.
- Krist S (2011). Volatile Compounds. The utilisation of volatile compounds in the characterisation of vegetable oils and fats and in reducing the bacterial count of ambient air. Wien: Peter Lang Europäischer Verlag der Wissenschaften.
- Lambert RJW et al. (2001). A study of the minimum inhibition concentration and mode of action of oregano essential oil, thymol and carvacrol. *Journal of Applied Microbiology* 91: 453–462.
- Longaray Delamare AP et al. (2007). Antibacterial activity of the essential oils of *Salvia officinalis* L. and *Salvia triloba* L. cultivated in South Brazil. *Food Chemistry* 100: 603–608.
- Malhi HK et al. (2017). Tea tree oil gel for mild to moderate acne; a 12 week uncontrolled, open-label phase II pilot study. *Australasian Journal of Dermatology* 58: 205–210.
- May J et al. (2000). Time-kill studies of tea tree oils on clinical isolations. *Journal of Antimicrobial Chemotherapy* 45: 639–643.
- Mondello F (2003). In vitro and in vivo activity of tea tree oil against azole-susceptible and -resistant human pathogenic yeasts. *Journal of Antimicrobial Chemotherapy* 51: 1223–1229.
- Mondello F et al. (2006). In vivo activity of terpinen-4-ol, the main bioactive component of *Melaleuca alternifolia* Cheel (tea tree) oil against azole-susceptible and -resistant human pathogenic *Candida* species. *BMC Infectious Diseases* 6: 158.
- Mulyaningsih S et al. (2010). Synergistic properties of the terpenoids aromadendrene and 1,8-cineole from the essential oil of *Eucalyptus globulus* against antibiotic-susceptible and antibiotic-resistant pathogens. *Phytomedicine* 17: 1061–1066.
- Nazzaro F et al. (2013). Effect of essential oils on pathogenic bacteria. *Pharmaceuticals* 6: 1451–1474.

- Nazzaro F et al. (2017). Essential oils and antifungal activity. *Pharmaceuticals* 10: 86.
- Ninomiya K et al. (2012). The essential oil of *Melaleuca alternifolia* (tea tree oil) and its main component, terpinen-4-ol protect mice from experimental oral candidiasis. *Biological and Pharmaceutical Bulletin* 35: 861–865.
- Noumi E et al. (2018). Chemical and biological evaluation of essential oils from cardamom species. *Molecules* 23: 2818.
- Nychas GJE (1995). Natural antimicrobials from plants. In: Gould GW. *New Methods of Food Preservation*. London: Blackie Academic & Professional; 58–89.
- Ohno T et al. (2003). Antimicrobial activity of essential oils against *Helicobacter pylori*. *Helicobacter* 8, 207–215.
- Oliveira JB et al. (2019). In vitro and in vivo antimicrobial activity of *Cymbopogon citratus* (DC.) Stapf. against *Staphylococcus* spp. isolated from newborn babies in an intensive care unit. *Microbial Drug Resistance* 25: 1490–1496.
- Özek G et al. (2010). Gas chromatographic-mass spectrometric analysis of volatiles obtained by four different techniques from *Salvia rosfolia* Sm., and evaluation for biological activity. *Journal of Chromatography A* 1217: 741–748.
- Pauli A (2001). Antimicrobial properties of essential oil constituents. *International Journal of Aromatherapy* 11: 126–133.
- Pauli A, Schilcher H (2004). Specific Selection of Essential Oil Compounds for Treatment of Children's Infection Diseases. *Pharmaceuticals* 1: 1–30.
- Pauli A, Schilcher H (2009). Anwendungsbeispiele mit dem Kamilleninhaltsstoff (-)-( $\alpha$ )-Bisabolol. *Infektionen, Entzündungen, Wunden: 18 Fallbeispiele zu (-)-( $\alpha$ )-Bisabolol. Komplementäre und integrative Medizin* 50: 28–32.
- Pauli A, Schilcher H (2016). In vitro antimicrobial activity of essential oils monographed in the European Pharmacopoeia 8th edition. In: Baser KHC, Buchbauer G. *Handbook of Essential Oils*. Boca Raton: Taylor & Francis Group; 431–611.
- Peixo LR et al. (2017). Antifungal activity, mode of action and anti-biofilm effects of *Laurus nobilis* Linnaeus essential oil against *Candida* spp. *Archives of Oral Biology* 73: 179–185.
- Pozzatti P et al. (2008). In vitro activity of essential oils extracted from plants used as spices against fluconazole-resistant and fluconazole-susceptible *Candida* spp. *Canadian Journal of Microbiology* 54: 950–956.
- Ratledge C, Wilkinson SG (1988). *An overview of microbial lipids*. In: Ratledge C, Wilkinson SG. *Microbial Lipids Vol. 1*. London: Academic Press Limited; 3–22.
- Reichling J (2017). Antibakterielle Wirkung ätherischer Öle unter besonderer Berücksichtigung multiresistenter Bakterien. *F-O-R-U-M* 49: 49–52.
- Reichling J (2020). Anti-biofilm and Virulence Factor-Reducing Activities of Essential Oils and Oil Components as a Possible Option for Bacterial Infection Control. *Planta Med* 86: 520–537.
- Reichling J et al. (2009). Essential oils of aromatic plants with antibacterial, antifungal, antiviral and cytotoxic properties - an overview. *Forschende Komplementärmedizin* 16: 79–90.
- Rosato A et al. (2009). In vitro synergic efficacy of the combination of Nystatin with the essential oils of *Origanum vulgare* and *Pelargonium graveolens* against some *Candida* species. *Phytomedicine* 16: 972–975.
- Satchell AC et al. (2002). Treatment of dandruff with 5% tea tree oil shampoo. *Journal of the American Academy of Dermatology* 47: 852–855.
- Schnitzler P et al. (2011). Chapter 10 - Antiviral effects of plant-derived essential oils and pure oil components. In: Thormar H. *Lipids and essential oils as antimicrobial agents*. John Chichester: Wiley & Sons; 239–254.
- Shaaban HAE et al. (2012). Bioactivity of essential oils and their volatile aroma compounds: Review. *Journal of Essential Oil Research* 24: 203–212.
- Shelef LA (1983). Antimicrobial effects of spices. *Journal of Food Safety* 6: 29–44.
- Sherry E (2001). Topical application of a new formulation of eucalyptus oil phytochemical clears methicillin-resistant *Staphylococcus aureus* infection. *American Journal of Infection Control* 29: 346.
- Stringaro A et al. (2014). Effects of *Mentha suaveolens* essential oil alone or in combination with other drugs in *Candida albicans*. Evidence-Based Complementary and Alternative Medicine: ArticleID 125904.
- Szweda P et al. (2015). Essential oils, silver nanoparticles and propolis as alternative agents against fluconazole resistant *Candida albicans*, *Candida glabrata* and *Candida krusei* clinical isolates. *Indian Journal of Microbiology* 55: 175–183.
- Utee A et al. (2002). The phenolic hydroxyl group of carvacrol is essential for action against the food-borne pathogen *Bacillus cereus*. *Applied Environmental Microbiology*: 1561–1568.
- Vivas R et al. (2019). Multidrug-resistant bacteria and alternative methods to control them: an overview. *Microbial Drug Resistance* 25: 890–908.
- Warnke PH et al. (2005). Antibacterial essential oils reduce tumor smell and inflammation in cancer patients. *Journal of Clinical Oncology* 23: 1588–1589.
- Weseler A (2005). A novel colorimetric broth microdilution method to determine the minimum inhibitory concentration (MIC) of antibiotics and essential oils against *Helicobacter pylori*. *Pharmazie* 60: 498–502.
- Zechner EL (2017). Relaxase and plasmid transfer in gram-negative bacteria. *Current Topics in Microbiology and Immunology* 413: 93–113.